
CO-OP PINNACLE BALANCER

GU	IARA	NTEE	D ANA	LYSIS

Crude Protein (min.)			28.00%
Lysine (min.)			
Methionine (min.)			0.60%
Crude Fat (min.)			3.00%
Crude Fiber (máx.)			8.00%
Acid Detergent Fiber (max.)			14.00%
Neutral Detergent Fiber (max.)			25.00%
Calcium (min.) Phosphorus (min.) Magnesium (min.)	2.50%	(max.)	3.50%
Phosphorus (min.)			1.50%
Magnesium (min.)			0.45%
Copper (min.)			200 ppm
Selenium (min.)			2 ppm
Zinc (min.)			500 ppm
Vitamin A'(min.)			24,000 ĺÚ/lb
Vitamin D (min.)			3,000 IU/lb
Vitamin E (min.)			500 IU/lb
Biotin (min.)			7.5 mg/lb
, ,	INGE	PEDIENTS	•



Dehulled Soybean Meal, Wheat Middlings, Corn Distillers Dried Grains with Solubles, Calcium Carbonate, Calcium Dehulled Soybean Meal, Wheat Middlings, Corn Distillers Dried Grains with Solubles, Calcium Carbonate, Calcium Phosphate, Molasses Products, Propionic Acid (a preservative), Flax Meal, Lysine, Salt, Lignin Sulfonate (pelleting agent), Magnesium Oxide, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Selenium Yeast, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, Threonine, Artificial Flavor, Vitamin E Supplement, Ascorbic Acid, Choline Chloride, Biotin, d-Calcium Pantothenate, Vitamin A Supplement, Vitamin D-3 Supplement, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Zinc Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Sodium Selenite, Cobalt Carbonate, Calcium Iodate, Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Sodium Bentonite, Calcium Bentonite.

FEEDING DIRECTIONS: Co-op Pinnacle Balancer is designed to be fed alone to fortify a forage-only diet. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7- to 10-day transitional period. Changes in the amount of a grain ration should not exceed 1.0 to 1.5 lb per day. Provide free-choice access to clean, fresh water and salt at all times. Store feed in a dry area protected from insects and rodents.

FEEDING GUIDLINES:

	Feeding Rate*			
Class	Ponies (600 to 900 lb)	Average Horses (900 to 1200 lb)	Large Horses (1200 to 1500 lb)	
Mature, Idle	0.5 to 1.0 lb	1.0 to 1.5 lb	1.5 to 2.0 lb	
Mature, Light Work	1.0 to 1.5 lb	1.5 to 2.0 lb	2.0 to 2.5 lb	
Mature, Moderate Work	1.5 to 2.0 lb	2.0 to 2.5 lb	2.5 to 3.0 lb	
Broodmare, Early/Mid Gestation	1.0 to 1.5 lb	1.5 to 2.0 lb	2.0 to 2.5 lb	
Broodmare, Late Gestation	1.5 to 2.0 lb	2.0 to 2.5 lb	2.5 to 3.0 lb	
Weanling	1.5 to 2.0 lb	2.0 to 2.5 lb	2.0 to 2.5 lb	
Yearling	1.0 to 1.5 lb	1.5 to 2.0 lb	2.0 to 2.5 lb	

Feeding rates are expressed as total lb of Pinnacle Balancer per head per day. All calculations assume a daily minimum intake of 1.5 lb of good-quality forage per 100 lb of body weight. Feeding amounts should be adjusted proportionately for horses with an ideal mature body weight of under 600 lb or over 1500 lb. Horses in heavy work, lactating broodmares and other horses requiring more calories than supplied by the feeding rates above should be switched to a more calorie-dense Pinnacle feed product designed to be fed at higher levels.

Manufactured By

Manufactured By PROTRITION FEED, LAVERGNE, TENNESSÉE 37086

ITEM NO. 336 TAG CODE NO. 102324-336

> Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.