CO-OP 36% TEND-R-LEEN BEEF CONCENTRATE PELLET-MON **MEDICATED**

For Growing Beef Steers and Heifers Fed in Confinement

Improved feed efficiency.

ACTIVE DRUG INGREDIENT

GUARANTEED ANALYSIS

GUARANTEED ANALTSIS			ıo
Crude Protein (min.)			
(This includes not more than 12.00% equivalent crude protein from non-protein nitrogen.)			
Crude Fat (min.)			
Crude Fiber (max.)			10.00%
Calcium (min.)	3.00%	(max.)	4.00%
Phosphorus (min.)			
Salt (min.)	2.50%	(max.)	3.50%
Potassium (min.)			
Vitamin A (min.)			
Vitamin E (min.)			

INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Calcium Carbonate, Urea, Salt, Molasses Products, Calcium Phosphate, Lignin Sulfonate, Animal Fat and Vegetable Oil with Ethoxyquin (a preservative), Vitamin E Supplement, Sodium Selenite, Magnesium Sulfate, Potassium Sulfate, Zinc Oxide, Calcium Iodate, Cobalt Carbonate, Manganous Oxide, Ferrous Oxide, Vitamin A Acetate (in gelatin), D-Activated Animal Sterol (source of Vitamin D-3), a-Tocopherol Acetate (source of Vitamin E), Choline Chloride, Vitamin B-12 Supplement, Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Thiamine Mononitrate, Pyridoxine Hydrochloride, Biotin, Folic Acid, Mineral Oil, Magnesium Oxide, Copper Sulfate, Zinc Sulfate, Manganese Sulfate, Ferrous Sulfate, Ethylenediamine Dihydriodide.

FEEDING DIRECTIONS: Feed continuously at the rate of 1.5 lb per head per day thoroughly mixed into the amount of shelled corn to be consumed daily to provide 180 mg of Monensin per head per day the provide 180 mg of Monensin per head per day the per head per day the per head per head

day. Use this ration after cattle have completed the Tend-R-Leen Starter Program. Always provide plenty of clean, fresh water.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in

calves to be processed for yeal.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. USE AS DIRECTED. Do not feed to sheep or other copper-sensitive animals.

Manufactured By PROTRITION FEED, LLC **LAVERGNE, TENNESSÉE 37086**

ITEM NO. 441 TAG CODE NO. 101223-441

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE