CO-OP 60% BEEF CONCENTRATE-MON

MEDICATED

For Growing Beef Steers and Heifers on Pasture (Stocker, Feeder and Slaughter) or in a Dry Lot; **Replacement Beef Heifers**

For increased rate of weight gain.

ACTIVE DRUG INGREDIENT

...... 304 g/ton Monensin

GUARANTEED ANALYSIS

Crude Protein (min.)			60.00%	
(This includes not more than 30.00% equivalent crude protein from non-protein nitrogen.)				
Crude Fat (min.)			0.50%	
Crude Fiber (max.)			10.00%	
Calcium (min.)	5.00%	(max.)	6.00%	
Phosphorus (min.)			0.60%	
Salt (min.)				
Potassium (min.)				
Zinc (min.)				
Vitamin A (min.)				
Vitamin E (min.)				



INGREDIENTS

Plant Protein Products, Calcium Carbonate, Urea, Salt, Molasses Products, Lignin Sulfonate, Calcium Phosphate, Vitamin A Supplement, Vitamin D-3 Supplement, Copper Sulfate, Vegetable Oil with Tertiary Butyl Hydroquinone (a preservative), Vitamin E Supplement, Sodium Selenite, Magnesium Oxide, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate. **FEEDING DIRECTIONS:** Co-op 60% Beef Concentrate-MON is to be fed at a rate to provide 50 to pro

200 mg of Monensin per head per day in not less than 1 lb of feed. During the first 5 days of feeding, cattle should receive 50 to 100 mg of Monensin per head per day in not less than 1 lb of feed. Use the mixing chart below to blend Co-op 60% Beef Concentrate-MON with corn. Feed at the rates indicated in the left column to provide 50 to 200 mg of Monensin per head per day. Be sure adequate pasture is available.

LB OF FEED	LB OF 60% BEEF	LB OF	PROTEIN
HEAD/DAY	CONCENTRATE-MON	CORN	<u>% OF MIX</u>
6.6 - 26.3	100	1900	11%
4.4 - 17.5	150	1850	12%
3.0 - 11.6	225	1775	14%
2.2 - 8.7	300	1700	16%

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do use in calves

WARNING: A withdrawal time has not been established for pre-ruminating carves. Do use in carves to be processed for veal. CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If eder feusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. USE AS DIRECTED. Do not feed to sheep or other conser-sensitive animals. or other copper-sensitive animals.

Manufactured By PROTRITION FEED, LLC LAVERGNE, TENNESSEE 37086

ITEM NO. 482 TAG CODE NO. 101223-482

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.