

# CO-OP CALF PRIMER/TCR I-MON

## MEDICATED

Starter for Growing Calves

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

### ACTIVE DRUG INGREDIENT

Monensin ..... 32 g/ton

### GUARANTEED ANALYSIS

Crude Protein (min.) .....	18.00%
Crude Fat (min.) .....	2.00%
Crude Fiber (max.) .....	15.00%
Acid Detergent Fiber (max.) .....	21.00%
Calcium (min.) ..... 0.70% (max.) .....	1.20%
Phosphorus (min.) .....	0.55%
Salt (min.) ..... 0.35% (max.) .....	0.85%
Potassium (min.) .....	0.80%
Selenium (min.) .....	0.3 ppm
Vitamin A (min.) .....	13,000 IU/lb

### INGREDIENTS

Processed Grain By-Products, Roughage Products, Plant Protein Products, Lignin Sulfonate, Molasses Products, Calcium Carbonate, Grain Products, Salt, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, Potassium Sulfate, Magnesium Sulfate, Calcium Phosphate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Selenium Yeast, Sodium Selenite, Magnesium Oxide, Copper Sulfate, Zinc Oxide, Zinc Sulfate, Manganese Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate.

**FEEDING DIRECTIONS:** Offer Co-op Calf Primer/TCR I-MON at the rate of 0.9 lb per 100 lb of body weight per day to attain the minimum level of Monensin intake for coccidiosis prevention. Intake of Monensin should not exceed 200 mg (12.5 lb of Co-op Calf Primer/TCR I-MON) per head per day. Always provide plenty of clean, fresh water. Do not let animals run out of feed.

**WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other copper-sensitive animals.

Manufactured By  
PROTRITION FEED, LLC  
LAVERGNE, TENNESSEE 37086

ITEM NO. 93115  
TAG CODE NO. 110524-93115

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

93115