

CO-OP CALF PRIMER/TCR II-MON

MEDICATED

Grower for Growing Calves

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Monensin 32 g/ton

GUARANTEED ANALYSIS

Crude Protein (min.)	14.00%
Crude Fat (min.)	1.50%
Crude Fiber (max.)	25.00%
Acid Detergent Fiber (max.)	29.50%
Calcium (min.) 0.60% (max.)	1.10%
Phosphorus (min.)	0.50%
Salt (min.) 0.40% (max.)	0.90%
Potassium (min.)	0.80%
Selenium (min.)	0.3 ppm
Vitamin A (min.)	9,000 IU/lb

INGREDIENTS

Roughage Products, Processed Grain By-Products, Plant Protein Products, Grain Products, Molasses Products, Calcium Carbonate, Lignin Sulfonate, Salt, Potassium Sulfate, Magnesium Sulfate, Calcium Phosphate, Vitamin A Supplement, Vitamin D-3 Supplement, Selenium Yeast, Sodium Selenite, Vitamin E Supplement, Magnesium Oxide, Copper Sulfate, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate.

FEEDING DIRECTIONS: Offer Co-op Calf Primer/TCR II-MON to calves weighing approximately 300 to 600 lb at the rate of 0.9 lb per 100 lb of body weight per day to attain the minimum level of Monensin intake for coccidiosis prevention. Intake of Monensin should not exceed 200 mg (12.5 lb of Co-op Calf Primer/TCR II-MON) per head per day. Always provide plenty of clean, fresh water. Do not let animals run out of feed.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other copper-sensitive animals.

Manufactured By
PROTRITION FEED, LLC
LAVERGNE, TENNESSEE 37086

ITEM NO. 93125

TAG CODE NO. 110524-93125

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

93125