

# CO-OP 14% PELLETTED BEEF DEVELOPER-MON MEDICATED

For Weaned Beef Cattle on Pasture

For increased rate of weight gain in growing cattle on pasture or in dry lot (stocker and feeder; beef replacement heifers).

### ACTIVE DRUG INGREDIENT

Monensin ..... 36 g/ton

### GUARANTEED ANALYSIS

Crude Protein (min.) .....		14.00%
Crude Fat (min.) .....		3.00%
Crude Fiber (max.) .....		20.00%
Calcium (min.) ..... 0.80%	(max.) .....	1.30%
Phosphorus (min.) .....		0.45%
Salt (min.) ..... 0.85%	(max.) .....	1.35%
Potassium (min.) .....		0.80%
Vitamin A (min.) .....		3,000 IU/lb

### INGREDIENTS

Processed Grain By-Products, Roughage Products, Plant Protein Products, Molasses Products, Propionic Acid (a preservative), Calcium Carbonate, Salt, Animal Fat and Vegetable Oil with Ethoxyquin (a preservative), Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Selenium Yeast, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Magnesium Oxide, Copper Sulfate, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate.

**FEEDING DIRECTIONS:** During the first 5 days of feeding, cattle should receive 2.8 to 5.5 lb per head per day of Co-op 14% Pelleted Beef Developer-MON. Following this adjustment period, feed at the rate of 2.8 to 11.1 lb per head per day to provide not less than 50 nor more than 200 mg of Monensin per head per day while having unlimited access to good-quality forage. Keep clean, fresh water available at all times.

**WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other copper-sensitive animals.

**Manufactured By  
PROTRITION FEED, LLC  
LAVERGNE, TENNESSEE 37086**

ITEM NO. 94176  
TAG CODE NO. 110722-94176

94176

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

**NET WEIGHT 50 lb (22.68 kg)/INVOICE**