

MARBLE LOCK BEEF FINISHER-MON

MEDICATED

For Growing Beef Steers and Heifers Fed in Confinement

For improved feed efficiency.

ACTIVE DRUG INGREDIENT

Monensin 30 g/ton

GUARANTEED ANALYSIS

Crude Protein (min.) 10.50%
Crude Fat (min.) 2.50%
Crude Fiber (max.) 15.00%
Calcium (min.) 0.35% (max.) 0.85%
Phosphorus (min.) 0.30%
Salt (min.) 0.15% (max.) 0.65%
Potassium (min.) 0.50%
Vitamin A (min.) 400 IU/lb

INGREDIENTS

Grain Products, Roughage Products, Processed Grain By-Products, Molasses Products, Plant Protein Products, Calcium Carbonate, Vegetable Oil with Tertiary Butyl Hydroquinone (a preservative), Salt, Calcium Phosphate, Lignin Sulfonate, Lysine, Magnesium Oxide, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Selenium Yeast, dl-Methionine Hydroxy Analogue Calcium, Calcium Hydroxide, Threonine, Vitamin E Supplement, Artificial Flavor, Biotin, Ascorbic Acid, Choline Chloride, d-Calcium Pantothenate, Vitamin A Supplement, Vitamin D-3 Supplement, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Zinc Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Sodium Selenite, Cobalt Carbonate, Calcium Iodate, Propionic Acid (a preservative), Acetic Acid, Sodium Propionate (a preservative), Ammonium Hydroxide, Sorbic Acid (a preservative), Magnesium Aluminosilicate, Sodium Bentonite, Calcium Bentonite.

FEEDING DIRECTIONS: Acclimate cattle to full feed using a high fiber grower feed medicated with Monensin. Then feed Marble Lock Beef Finisher-MON continuously at the rate of 4 to 32 lb per head per day to provide 60 to 480 mg of Monensin per head per day. Always provide plenty of clean, fresh water. Keep feed trough clean and free of old or out of condition feed and feed fines. Consult local feed representative for specific ration balancing.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other copper-sensitive animals.

Manufactured By
PROTRITION FEED, LLC
LAVERGNE, TENNESSEE 37086

ITEM NO. 94471

TAG CODE NO. 052924-94471

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

94471