CO-OP SUPREME CATTLE MINERAL-MON

MEDICATED For Beef Cows on Pasture

For improved feed efficiency.

ACTIVE DRUG INGREDIENT Monensin..... 1,200 g/ton

GUARANTEED ANALYSIS

Calcium (min.)	16.50%	(max.)	19.50%
Phosphorus (min.)		·····	4.00%
Salt (min.)	20.00%	(max.)	24.00%
Magnesium (min.)			3.50%
Copper (min.)			1,950 ppm
Manganese (min.)			3,000 ppm
Selenium (min.)			30 ppm
Zinc (min.)			3,950 ppm
Vitamin A (min.)			260,000 IU/lb
Vitamin D-3 (min.)			60,000 IU/lb
Vitamin E (min.)			450 IU/lb

INGREDIENTS

Calcium Carbonate, Salt, Calcium Phosphate, Magnesium Oxide, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Bentonite, Molasses Products, Yeast Extract, Yeast Culture, Vitamin A Supplement, Vitamin D-3 Supplement, Manganese Hydroxychloride, Selenium Yeast, Iron Oxide, Basic Copper Chloride, Ethylenediamine Dihydriodide, Vitamin E Supplement, Natural and Artificial Flavors, Zinc Hydroxychloride, Sodium Selenite.

FEEDING DIRECTIONS: Hand feed at the rate of 100 g (3.53 oz) per head per day in not less than 1 lb of feed to beef cows on pasture to provide 132 mg of Monensin and 3 mg of selenium per head per day. During the first five days, cattle should receive no more than 100 mg of Monensin (2.66 oz of Co-op Supreme Cattle Mineral-MON) per head per day. This feeding rate is Zinpro® verified to provide 360 mg of Zn and 12.6 mg of Co from Zinpro® products. Provide plenty of good-quality forage and clean, fresh water continuously.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other coppersensitive animals.

> **Manufactured By** PROTRITION FEED, LLC LAVERGNE, TENNESSEE 37086

ITEM NO. 96561 TAG CODE NO. 102224-96561

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE