



BEEF PELLETS-MON

MEDICATED

For Weaned Beef Cattle

For increased rate of weight gain in growing beef steers and heifers on pasture (stocker, feeder and slaughter) or in a dry lot; replacement beef heifers. For improved feed efficiency in growing beef steers and heifers fed in confinement.

ACTIVE DRUG INGREDIENT

Monensin 27 g/ton

GUARANTEED ANALYSIS

| | |
|---|-------------|
| Crude Protein (min.) | 13.00% |
| Crude Fat (min.) | 2.50% |
| Crude Fiber (max.) | 22.50% |
| Calcium (min.) 1.50% (max.) | 2.00% |
| Phosphorus (min.) | 0.40% |
| Salt (min.) 0.40% (max.) | 0.90% |
| Potassium (min.) | 0.65% |
| Vitamin A (min.) | 2,000 IU/lb |

INGREDIENTS

Roughage Products, Processed Grain By-Products, Plant Protein Products, Calcium Carbonate, Molasses Products, Propionic Acid (a preservative), Grain Products, Salt, Animal Fat and Vegetable Oil with Ethoxyquin (a preservative), Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Magnesium Oxide, Copper Sulfate, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate.

FEEDING DIRECTIONS FOR PASTURE STEERS AND HEIFERS: During the first 5 days of feeding, cattle should receive 3.7 to 7.4 lb per head per day of Cumberland Mills Beef Pellets-MON. Following this adjustment period, feed at the rate of 3.7 to 14.8 lb per head per day to provide not less than 50 nor more than 200 mg of Monensin per head per day while on pasture. Keep clean, fresh water available at all times.

FEEDING DIRECTIONS FOR STEERS AND HEIFERS FED IN CONFINEMENT: Feed continuously at the rate of 3.7 to 35.5 lb per head per day to provide 50 to 480 mg of Monensin per head per day along with adequate roughage. Keep clean, fresh water available at all times.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other copper sensitive animals.

Manufactured By
PROTRITION FEED, LLC
LAVERGNE, TENNESSEE 37086

ITEM NO. CM401
TAG CODE NO. 050423-CM401

Seller warrants only that this feed conforms to the description on the label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

CM401